



A, B: 山田和彦 (三浦半島)

日本近海 Japanese Waters

ML 65mm. 外套膜は卵楕円形。外套幅は外套長の45%。背側は紫褐色で赤斑がある。鰭の後端が耳型につまみ出したように張り出す。腹部には鰭に沿って赤褐色の縦線がある。I腕は他腕より長く雄では先端の1/3の保護膜が広がる。触腕吸盤は不等大で、中央の4~5個が大きい。貝殻は *Doratosepion* 型で、長さは幅の約4倍。本州中部以西の陸棚帯。

Small sepiid. MW 45% of ML. Dorsal side brownish purple, scattered by red spots. Posterior end of fin slightly expanded like an "ear". Reddish orange stripe running along ventral fin base. Arm I longer than others, distal 1/3 in male bearing frilled protective membrane. Tentacular club suckers subequal, with large 4-5 medial ones. Shell *Doratosepion*-type, SL four times of SW. *S. (D.) misakiensis* differs from this species in having slender body (MW less than 30% of ML) and fused "ears". Shelf in the west of middle Honshu.

[A. 背面 (♂) : B. 腹面 (♂) : C. 触腕]



A: 窪寺恒己 (島根県). B, C: Sasaki (1929)

日本近海 Japanese Waters

ML 65mm. スジコウイカに似るがやや細い。外套膜の腹側には鰭に沿って赤褐色の縦帯が走る。鰭の後端はやや耳状になるが、基部では左右片葉は癒合している。I腕は外套長の半分より短い。触腕掌部は半月型。吸盤は8列で中央の数個は僅かに大きい。貝殻は *Doratosepion* 型で、長さは幅の4倍。内円錐は低い。相模湾、駿河湾、日本海西部の陸棚。

Resembling *S. tokioensis*, but slightly slenderer. Brownish red stripes running along fin base on ventral mantle. Posterior end of fins lobed, but both lobes fused together basally. Arm I shorter than half of ML. Tentacular club crescent in shape, with eight rows of suckers of which mesial several ones are slightly enlarged. Shell *Doratosepion*-type, with length four times width. Inner cone low and flat. Sagami Bay, Suruga Bay and western Japan Sea.

[A. 背面 (60mm) : B. 背面 (♂60mm) : C. 触腕]

